

September 2017 Sea Ice Outlook

(June Issue)

Same report used for July

Canadian Ice Service

Environment Canada's Canadian Ice Service (CIS) is predicting the 2017 minimum Arctic sea extent at $4.0 \times 10^6 \text{ km}^2$.

As with previous CIS contributions, the 2017 forecast was derived by considering a combination of methods: 1) a qualitative heuristic method based on observed end-of-winter Arctic ice thickness/extent, as well as winter surface air temperature, spring ice conditions and the summer temperature forecast; 2) two simple statistical methods based on an Optimal Filtering Based Model (OFBM), that uses an optimal linear data filter to extrapolate the September sea ice extent time-series into the future and 3) a Multiple Linear Regression (MLR) prediction system that tests ocean, atmosphere and sea ice predictors.

Based on winter air temperatures and sea ice extents and thickness, a September 2015 minimum ice extent value of $3.8 \times 10^6 \text{ km}^2$ is heuristically predicted. The CIS OFB models predict $4.24 \times 10^6 \text{ km}^2$ and $3.78 \times 10^6 \text{ km}^2$ and the CIS MLR model predicts $4.2 \times 10^6 \text{ km}^2$. The average forecast value of the four methods combined is $4.0 \times 10^6 \text{ km}^2$.

Heuristic Forecast

The CIS heuristic for the 2017 September minimum is $3.8 \times 10^6 \text{ km}^2$. The uncertainty in the heuristic forecast is estimated at $\pm 0.2 \times 10^6 \text{ km}^2$

Rational:

Winter conditions (Nov-Apr)


- Above normal air temperatures across most of the Canadian Arctic and very much higher than normal temperatures over the Laptev and Chukchi Seas as well as most of the Arctic Basin. Labrador saw near normal temperatures
- At the beginning of the winter significantly less MYI than normal in the Beaufort Sea. Normally the Beaufort Sea area is covered with almost 52% MYI however this year only 2.3% was present at the beginning of January 2017.
- Many regions of the northern Canadian Arctic Archipelago also had less old ice than normal.
- However the old ice from the northern Canadian Arctic Archipelago percolated southwards into the southern part of the archipelago due to the fracturing and melting of the ice. It permitted the flow of old ice further south.
- *April ice thickness from Cryosat shows thicker ice north of the Canadian Archipelago compared to the record 2012*
- Measured fast ice thicknesses at select stations throughout the Canadian Arctic were near normal to below normal at the end of April

Spring ice conditions

- Early spring melt in the southern Beaufort Sea and Baffin Bay as well as northwestern and eastern Hudson Bay
- Fast ice fracture and melt season is up to 3 weeks ahead of normal throughout the CAA

Summer temperature forecast

- Above normal temperatures forecast through to October over the entire Arctic basin except for over the Davis Strait /Labrador Sea where forecast is for slightly below normal temperatures.



STATISTICS BASED UPON 1981-2010
LES STATISTIQUES BASÉES SUR 1981-2010

Figure 1. January 2, 2017 anomalies in multi-year ice concentrations (1981-2010 climatology)

Statistical Method #1: Optimal Filtering Based Model

Forecast: The 2017 forecast for the September sea ice extent is **4.24*10⁶ km²**.

Model Details/References

- Details of the Optimal Filtering Based Model (OBFM) used here, as well as model code, can be found in: Press, W.H., S.A. Teukolsky, W.T. Vetterling and B.P. Flannery (1992): Numerical Recipes in Fortran 77, Second Edition: The art of scientific computing. Cambridge University Press, Cambridge UK [Chapter 13, section 13.6]
- Models based on optimal linear data filters have proven skill at predicting other climate indices (e.g. Nino3 and Nino3.4 SSTs): 1) Kim, K-Y., and G.R. North (1998): EOF-Based Linear Prediction Algorithm: Theory. J Clim, 11, 3045-3056. 2) Kim, K-Y, and G.R. North (1999): EOF-Based Linear Prediction Algorithm: Examples. J.Clim, 12, 2076-2092.

Statistical Method #2: Andre

Forecast: The 2017 forecast for the September sea ice extent is **3.78*10⁶ km²**.

Model Details/References

April, A. , 2017. Statistical Forecast Model for Ice-Related Events in the Arctic, Weather and Forecasting, 32, 469-478, DOI: 10.1175/WAF-D-16-0139.1.

Statistical Method #3: Multiple Linear Regression Model

Forecast: The 2017 forecast for the September sea ice extent is **4.2 *10⁶ km²**.

Model Details/References

The regression model was generated using an automated selection scheme (Tivy et al. 2007) that uses step-wise regression and limits the number of predictors to only 2. Predictors included in the original predictor pool: pan-arctic (60N-90N) SLP and SAT; northern hemisphere z500, global SST; monthly atmosphere teleconnection indices from NOAA/CPC, monthly AO, monthly SOI, monthly PDO and monthly sea ice extent.

Tivy, A., B.Alt, S.E.L. Howell, K. Wilson and J.J. Yackel (2007). Long-range prediction of the shipping season in Hudson Bay: A statistical approach. Weather and Forecasting, 22, 1063-1075, doi:10.1175/WAF1038.WAF10

Forecast Verification: Past June Outlooks vs 3 Benchmark Models

| Year | Observed | CIS Forecast | | Heuristic Forecast | | OFB Forecast | | MLR Forecast | |
|------|----------|--------------|-------|--------------------|-------|--------------|-------|--------------|-------|
| | | Forecast | Diff. | Forecast | Diff. | Forecast | Diff. | Forecast | Diff. |
| 2009 | 5.66 | 5 | -0.66 | 5 | -0.66 | 4.2 | -1.16 | 5.65 | 0.29 |
| 2010 | 4.9 | 4.85 | -0.05 | 4.85 | -0.05 | 4.91 | 0.01 | 5.7 | 0.8 |
| 2011 | 4.61 | 4.9 | 0.29 | 4.7 | 0.09 | 4.8 | 0.19 | 5.6 | 0.99 |
| 2012 | 3.61 | 4.7 | 1.09 | 4.75 | 1.14 | 4.3 | 0.69 | 5.1 | 1.49 |
| 2013 | 5.35 | 3.8 | -1.55 | 3.6 | -1.75 | 4.05 | -1.3 | 5 | -0.35 |
| 2014 | 5.02 | 4.9 | -0.12 | 4.8 | -0.22 | 4.37 | -0.65 | 5.6 | 0.58 |
| 2015 | 4.63 | 4.7 | 0.07 | 4.5 | -0.13 | 4.42 | -0.21 | 5.3 | 0.67 |
| 2016 | 4.14 | 4.3 | 0.16 | 4.3 | 0.16 | 4.1 | -0.04 | 4.6 | 0.46 |
| 2017 | | 4.0 | | 3.8 | | 4.24 | | 4.2 | |
| MAE | | | 0.50 | | 0.53 | | 0.53 | | 0.7 |

Table 1. Verification of the CIS June outlook submissions starting in 2009. The final CIS forecast is a combination of a heuristic forecast and two statistical methods, optimal filtering based model and a multiple linear regression model.

| Year | Observed | Climatology Forecast | | Trend Forecast | | Persistence Forecast | |
|------|----------|----------------------|-------|----------------|-------|----------------------|-------|
| | | Forecast | Diff. | Forecast | Diff. | Forecast | Diff. |
| 2009 | 5.66 | 6.32 | 0.93 | 5.25 | -0.14 | 6.44 | 1.05 |
| 2010 | 4.9 | 6.24 | 1.31 | 5.15 | 0.22 | 6.12 | 1.19 |
| 2011 | 4.61 | 6.17 | 1.54 | 4.89 | 0.26 | 5.8 | 1.17 |
| 2012 | 3.61 | 6.07 | 2.46 | 4.63 | 1.02 | 5.98 | 2.37 |
| 2013 | 5.35 | 5.88 | 0.53 | 4.31 | -1.04 | 5.76 | 0.41 |
| 2014 | 5.02 | 5.82 | 0.8 | 4.27 | -0.75 | 5.4 | 0.38 |
| 2015 | 4.63 | 5.71 | 1.08 | 4.27 | -0.36 | 5.28 | 0.65 |
| 2016 | 4.14 | 5.63 | 1.19 | 4.06 | -0.08 | 4.75 | 0.61 |
| 2017 | | 5.58 | | 4.02 | | 5.25 | |
| MAE | | | | | | | |

Table 2. Verification of 3 benchmark models for the June outlook submission starting in 2009. The most common benchmark models are climatology, extrapolation of the long-term trend and anomaly persistence (May). The benchmark model forecasts are based on the previous 20 years as the training period. For example, the 2009 forecast is based on the 1989 – 2008 training period and 2014 is based on the 1993-2014 training period.