

Sea Ice Outlook
2020 August Report
Individual Outlook

Name of contributor or name of contributing organization:

McGill Team

Is this contribution from a person or group not affiliated with a research organization?

Name and organization for all contributors. Indicate primary contact and total number of people who may have contributed to your Outlook, even if not included on the author list.

Bruno Tremblay (1), Erik Johnson (2), Charles Brunette (1)

1) McGill University, Montreal, Canada

2) University of Otago, Dunedin, New Zealand

**Do you want your June contribution to automatically be included in subsequent reports?
(If yes, you may still update your contribution via the submission form.)**

Include this submission in both the July and August reports.

What is the type of your Outlook projection?

Statistical

Starting in 2017 we are accepting both pan-Arctic and pan-Antarctic sea ice extent (either one or both) of the September monthly mean. As in 2016, we are also collecting Alaskan regional sea ice extent. To be consistent with the validating sea ice extent index from NSIDC, if possible, please first compute the average sea ice concentration for the month and then compute the extent as the sum of cell areas > 15%.

a) Pan-Arctic September extent prediction in million square kilometers.

3.89

b) same as in (a) but for pan-Antarctic. If your method differs substantially from that for the Arctic, please enter it as a separate submission.

c) same as in (b) but for the Alaskan region. Please also tell us maximum possible extent if every ocean cell in your region were ice covered.

"Executive summary" of your Outlook contribution (using 300 words or less) describe how and why your contribution was formulated. To the extent possible, use non-technical language.

Our research focuses on seasonal predictability of sea ice in the Arctic Ocean, using observations-based approaches. We are interested in the winter preconditioning effect on the pack ice before the summer melt. Specifically, we investigate how dynamic processes affect preconditioning, in other words, we ask how anomalies in the general circulation of sea ice will influence later conditions of the Arctic Ocean pack ice under a typical melt season. We investigate the skill of different sea ice predictors, including atmospheric forcing parameters that physically connect to wintertime sea ice dynamics.

The dovekSIE method builds on the correlation between winter Fram Strait sea ice export and the following September minimum sea ice extent, presented in Williams et al., 2016. A positive anomaly of the winter Fram Strait sea ice export is associated with enhanced circulation of ice through the Transpolar Drift Stream and positive anomalies of coastal divergence of sea ice along the Eurasian coastlines. Increased coastal divergence late in the winter causes anomalies of younger and thinner ice in the peripheral seas, which is more vulnerable to melting in the summer.

The dovekSIE forecasts are generated using the sea level pressure difference between Greenland and Svalbard as a proxy for area of ice exported through Fram Strait. Sea ice tends to flow parallel to isobars and the pressure difference across Fram Strait correlates with sea ice export ($r=0.44$). Sea level pressure fields are available in near-real-time and therefore enable the continuous update of dovekSIE forecasts during winter via the web app.

We are supporting the activities of the Sea Ice Prediction Network with great enthusiasm. This is our fourth contribution to the Sea Ice Outlook.

Brief explanation of Outlook method (using 300 words or less).

regional SIE is 0.90 (correlation using cross-validated experiments) with RMSE of 0.22 million

Tell us the dataset used for your initial Sea Ice Concentration (SIC).

Sea ice concentration is not used as an initial condition (such as in a dynamical model). However, we use sea ice extent from the NSIDC Sea Ice Index V3 to fit our statistical model. <https://doi.org/10.7265/N5K072F8>

Tell us the dataset used for your initial Sea Ice Thickness (SIT) used. Include name and date.

If you use a dynamic model, please specify the name of the model as a whole and each component including version numbers and how the component is initialized:

[DynamicModelType]

If available from your method.

a) Uncertainty/probability estimates:

Median

Ranges

Standard Deviations

b) Brief explanation/assessment of basis for the uncertainty estimate (1-2 sentences).

RMSE = 0.46 million km². We compare hindcasts to the observed mean September sea ice extent for the 1993-2019 period.

c) Brief description of any post processing you have done (1-2 sentences).