

## Walt Meier, NSIDC, August 2020 Sea Ice Outlook Contribution

Name: Walt Meier  
Organization: National Snow and Ice Data Center  
Label: Meier (NSIDC)  
Contact: walt@nsidc.org

**Executive Summary:** This method applies daily ice loss rates to extrapolate from the start date (August 1) through the end of September. Projected September daily extents are averaged to calculate the projected September average extent. Individual years from 2005 to 2017 are used, as well as averages over 1981-2010 and 2007-2019. The 2007-2019 average daily rates are used to estimate the official submitted estimate.

The predicted September average extent for 2020 is 3.93 ( $\pm 0.36$ ) million square kilometers, which is a substantial decrease from the July estimate of 4.44 million square kilometers. The minimum daily extent is predicted to be 3.81 ( $\pm 0.37$ ) million square kilometers, a decrease from July's 4.32 million square kilometers and occurs on 14 September. The range of estimates has decreased ( $\pm 0.36$ ) due to the shorter remainder of the melt season. Based on the last 15 years, two of the projections (2008 and 2012) yield an extent lower than the current record low September extent of 3.57 million sq km in 2012. Thus, there is a roughly a 2 in 15 (13%) of a new record September extent in the Arctic

Using the same method, the predicted Antarctic average extent for September 2020 is 18.09 ( $\pm 0.30$ ) million square kilometers. The maximum daily extent is predicted to be 18.20 ( $\pm 0.32$ ) million square kilometers and occurs on 30 September.

| Start Date | Arctic Sep  | Arctic Min  | Antarctic Sep | Antarctic Max |
|------------|-------------|-------------|---------------|---------------|
| 1 June     | 4.13 (0.66) | 4.01 (0.66) | 18.39 (0.61)  | 18.50 (0.63)  |
| 1 July     | 4.44 (0.53) | 4.32 (0.53) | 18.16 (0.57)  | 18.27 (0.58)  |
| 1 August   | 3.93 (0.36) | 3.81 (0.37) | 18.09 (0.30)  | 18.20 (0.32)  |

*Table 1.* Predicted 2020 September extent and daily minimum extent (in million sq km) based on extent loss rates for 2007-2019 and, in parenthesis, the standard deviation, for the Outlook contributions in June, July, and August.

Outlook type: Statistical

Initial SIC data set used: NASA Team algorithm extents from the NSIDC Sea Ice Index, Version 3 ([http://nsidc.org/data/seaice\\_index/](http://nsidc.org/data/seaice_index/)).

Initial SIT data used: N/A

Prediction of September pan-Arctic extent: 3.93 ( $\pm 0.36$ ) million square kilometers

Prediction of week that minimum daily extent will occur: Week of 9 September (specific day predicted is 14 September).

Outlook Description: This method applies daily ice loss rates to extrapolate from the start date (1 August) through the end of September. Projected September daily extents are averaged to calculate the projected September average extent. Individual years from 2007 to 2019 are used, as well as averages over 1981-2010 and 2007-2019. The 2007-2019 average daily rates are used to estimate the official submitted estimate. The method essentially provides the range of September extents that can be expected based on how the ice has declined in past years, though it is possible that record fast or slow daily loss rates may yield a value outside the projected range. It also can provide a probability of a new record by comparing how many years of loss rates yield a record relative to all years. It has the benefit that it can easily and frequently (daily if desired) be updated to provide updated estimates and probabilities and as the minimum approaches the “window” of possible outcomes narrows.

### **Detailed Description and Discussion**

This method is a simple statistical method that uses previous years’ daily rates of extent change to project the 2020 daily extent through the end of September. The monthly average is then calculated from the September daily extents. This year, the last twelve years (2007 – 2019) are used for the projection because these years are more representative of recent conditions than using all years in the 40-year time series.

This method, initialized with 1 August extent, yields a September 2020 extent of 3.93 ( $\pm 0.36$ ) million sq km. The range in values, calculated from the standard deviation of the twelve years, reflects the significant amount of variability in daily extent change rates between 1 August and 30 September. Using the standard 30-year 1981-2010 climatology, the September extent is 4.21 million sq km. The lowest projected extent (from the last 15 years), which is from the 2012 rates, is 3.21 million sq km, while the highest, from 2006 rates, is 4.50 million square kilometers. The daily extent trajectories for the two averages and the high and low years from the last decade are provided in the figure below and the prediction using the rates for each of the last ten years are provided in the table below.

The expected skill has increased over June and July because there is one or two months less of potential variability, so the range of estimates has decreased from  $\sim 0.53$  to 0.36 million sq km. Thus, by August, the method starts to provide a reasonable envelop of physically realistic September extents. As September approaches, the “window” of possible extents will continue to narrow and hones in on the final observed extent.

| Year                     | September Extent | Daily Minimum Extent |
|--------------------------|------------------|----------------------|
| 2005                     | 4.29             | 4.10                 |
| 2006                     | 4.50             | 4.40                 |
| 2007                     | 3.73             | 3.60                 |
| 2008                     | <b>3.38</b>      | <b>3.28</b>          |
| 2009                     | 4.29             | 4.16                 |
| 2010                     | 4.03             | 3.79                 |
| 2011                     | 3.94             | 3.74                 |
| 2012                     | <b>3.21</b>      | <b>3.05</b>          |
| 2013                     | 4.27             | 4.13                 |
| 2014                     | 4.19             | 4.00                 |
| 2015                     | 3.74             | 3.59                 |
| 2016                     | 3.79             | 3.48                 |
| 2017                     | 4.15             | 4.10                 |
| 2018                     | 4.10             | 3.96                 |
| 2019                     | 4.23             | 4.07                 |
| <i>Average 2007-2019</i> | 3.93             | 3.81                 |
| <i>Average 1981-2010</i> | 4.21             | 4.09                 |

Table 2. Predicted 2020 September extent and daily minimum extent using ice loss rates from different years or multi-year averages. Values in **bold** represent cases that would set new record lows.

Daily trajectory of sea ice extent from previous years' rates  
Arctic Meier (National Snow and Ice Data Center)





Figure 1. Predicted Arctic trajectories from the 1 August observed extent. The official estimate is based on the projection used the average of the last 15 years (2005-2019, green solid line) and for the 1981-2010 average (solid red line). The rates for 2006 and 2012 (dotted lines) yield the highest and lowest predicted extents respectively.

We use the same methodology for the Antarctic. However, since the trend has not particularly accelerated there isn't a reason to use any particular subset of years. It is clear from Figure 2 that there is very little difference between using the thirty-year average versus the 13-year (2007-2019) average employed for the Arctic. So, the projection for the Antarctic is based on the daily rates of change for the 2007-2019 average to be consistent with the Arctic. The highest project results from using 2006 rates and the lowest projection comes from 2015 rates.

Using the 2007-2019 rates, the projected Antarctic September average is 18.09 ( $\pm 0.30$ ) million square kilometers. Over the last 15 years, the highest September extent of 18.65 million square kilometers came from using 2006 rates, and the lowest September extent, using 2015 rates is 17.62 million square kilometers. The very large spread between the highest and lowest is not surprising given the high interannual variability in Antarctic sea ice. Since the Antarctic sea ice at its maximum encircles the entire continent over a thousand kilometers from the coast in most regions, even relatively small differences in ice edge location can result in large differences in total area. The maximum daily extent is predicted to be 18.20 ( $\pm 0.32$ ) million square kilometers and occurs on 30 September.



**Figure 2.** Predicted Antarctic trajectories from the 1 August observed extent. The official estimate is based on the projection used the average of the last 13 years (2007-2018, green solid line) and for the 1981-2010 average (solid red line). The rates for 2006 and 2015 (dotted lines) yield the highest and lowest predicted extents respectively.

### **Input Data Set References**

Maslanik, J. and J. Stroeve. 1999, updated daily. Near-Real-Time DMSP SSMIS Daily Polar Gridded Sea Ice Concentrations, Version 1. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center. doi: <https://doi.org/10.5067/U8C09DWVX9LM>.

Fetterer, F., K. Knowles, W. Meier, M. Savoie, and A. K. Windnagel. 2017, updated daily. Sea Ice Index, Version 3. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center. doi: <https://doi.org/10.7265/N5K072F8>.