## Sea Ice Outlook

## 2022 September Report Individual Outlook

Nan	ie of	contributor	or nan	ie of o	contributing	organization:

NSIDC (Meier)

Is this contribution from a person or group not affiliated with a research organization?

Name and organization for all contributors. Indicate primary contact and total number of people who may have contributed to your Outlook, even if not included on the author list.

NSIDC (Meier)

Do you want your June contribution to automatically be included in subsequent reports? (If yes, you may still update your contribution via the submission form.)

[Do you want your contribution for this month to automatically be included in subsequent reports?]

What is the type of your Outlook projection?

Statistical/ML

Starting in 2017 we are accepting both pan-Arctic and pan-Antarctic sea ice extent (either one or both) of the September monthly mean. As in 2016, we are also collecting Alaskan regional sea ice extent. To be consistent with the validating sea ice extent index from NSIDC, if possible, please first compute the average sea ice concentration for the month and then compute the extent as the sum of cell areas > 15%.

a) Pan-Arctic September extent prediction in million square kilometers.

b) same as in (a) but for pan-Antarctic. If your method differs substantially from that for the Arctic, please enter it as a separate submission.

17.7

c) same as in (b) but for the Alaskan region. Please also tell us maximum possible extent if every ocean cell in your region were ice covered.


"Executive summary" of your Outlook contribution (using 300 words or less) describe how and why your contribution was formulated. To the extent possible, use non-technical language.

This method applies daily ice loss rates to extrapolate from the start date (September 1) through the end of September. Projected September daily extents are averaged to calculate the projected September average extent. Individual years from 2005 to 2021 are used, as well as averages over 1981-2010 and 2007-2021. The 2007-2021 average daily rates are used to estimate the official submitted estimate.

The predicted September average extent for 2022 is  $5.03~(\pm0.09)$  million square kilometers. The minimum daily extent is predicted to be  $4.91~(\pm0.10)$  million square kilometers and occurs on 16 September. The range of estimates reflects the variability in ice loss rates over the final month of the melt season. Based on the last 17 years (2005-2021), there is a 0% chance that 2022 will be lower than the current record low September extent of 3.57 million sq km in 2012. Using the same method, the predicted Antarctic average extent for September 2022 is  $17.70~(\pm0.26)$  million square kilometers. The maximum daily extent is predicted to be  $17.79~(\pm0.31)$  million square kilometers and occurs on 26 September.

## Brief explanation of Outlook method (using 300 words or less).

This method applies daily ice loss rates to extrapolate from the start date (September 1) through the end of September. Projected September daily extents are averaged to calculate the projected September average extent. Individual years from 2005 to 2021 are used, as well as averages over 1981-2010 and 2007-2021. The 2007-2021 average daily rates are used to estimate the official submitted estimate. The method essentially provides the range of September extents that can be expected based on how the ice has declined in past years, though it is possible that record fast or slow daily loss rates may yield a value outside the projected range. It also can provide a probability of a new record by comparing how many years of loss rates yield a record relative to all years. It has the benefit that it can easily and frequently (daily if desired) be updated to



c) Brief description of any post-processing you have done (1-2 sentences).							
None							