

## Walt Meier, NSIDC, August 2023 Sea Ice Outlook Contribution

Name: Walt Meier  
Organization: National Snow and Ice Data Center  
Label: Meier (NSIDC)  
Contact: walt@colorado.edu

Executive Summary: This method applies daily ice loss rates to extrapolate from the start date (August 1) through the end of September. Projected September daily extents are averaged to calculate the projected September average extent. Individual years from 2005 to 2022 are used, as well as averages over 1981-2010 and 2007-2022. The 2007-2022 average daily rates are used to estimate the official submitted estimate.

The predicted September average extent for 2023 is 4.77 ( $\pm 0.35$ ) million square kilometers. The minimum daily extent is predicted to be 4.64 ( $\pm 0.35$ ) million square kilometers and occurs on 16 September. This is an increase of  $\sim 0.15$  million square kilometers from the July predictions. The range of the estimates has declined, reflecting the smaller potential variability in ice loss rates over the final  $\sim 1.5$  months of the melt season compared to June and July. None of the last 18 years (2005-2022) yield a 2023 extent lower than the current record low September extent of 3.57 million sq km in 2012.

Using the same method, the predicted Antarctic average extent for September 2023 is 16.22 ( $\pm 0.34$ ) million square kilometers. The maximum daily extent is predicted to be 16.31 ( $\pm 0.34$ ) million square kilometers and occurs on 26 September. These values are higher, by  $\sim 0.5$  million square kilometers, than the July predictions.

Outlook type: Statistical

Initial SIC data set used: NASA Team algorithm extents from the NSIDC Sea Ice Index, Version 3 ([http://nsidc.org/data/seaice\\_index/](http://nsidc.org/data/seaice_index/)).

Initial SIT data used: N/A

Prediction of September Pan-Arctic extent: 4.77 ( $\pm 0.35$ ) million square kilometers  
Prediction of September Pan-Antarctic extent: 16.22 ( $\pm 0.34$ ) million sq km

September pan-Arctic Anomaly Forecast: +0.58 million square kilometers (2023 trend extrapolation value = 4.19 million square kilometers)

Prediction of week that minimum daily extent will occur: Week of 12 September (specific day predicted is 16 September).

Outlook Description: This method applies daily ice loss rates to extrapolate from the start date (August 1) through the end of September. Projected September daily extents are averaged to calculate the projected September average extent. Individual

years from 2005 to 2022 are used, as well as averages over 1981-2010 and 2007-2022. The 2007-2022 average daily rates are used to estimate the official submitted estimate. The method essentially provides the range of September extents that can be expected based on how the ice has declined in past years, though it is possible that record fast or slow daily loss rates may yield a value outside the projected range. It also can provide a probability of a new record by comparing how many years of loss rates yield a record relative to all years. It has the benefit that it can easily and frequently (daily if desired) be updated to provide updated estimates and probabilities and as the minimum approaches the “window” of possible outcomes narrows.

### Detailed Description and Discussion


This method is a simple statistical method that uses previous years’ daily rates of extent change to project the 2023 daily extent through the end of September. The monthly average is then calculated from the September daily extents. This year, the last 16 years (2007 – 2022) are used for the projection because these years are more representative of recent conditions than using all years in the 45-year time series.

<b>Year</b>	<b>September Extent</b>	<b>Daily Minimum Extent</b>
2005	5.09	4.90
2006	5.30	5.20
2007	4.52	4.40
2008	4.18	4.08
2009	5.09	4.96
2010	4.83	4.59
2011	4.74	4.54
2012	4.01	3.84
2013	5.07	4.94
2014	4.99	4.80
2015	4.54	4.39
2016	4.59	4.28
2017	4.95	4.81
2018	4.90	4.76
2019	5.03	4.86
2020	4.81	4.64
2021	5.26	5.08
2022	4.81	4.61
<i>Average 2007-2022</i>	4.77	4.64
<i>Average 1981-2010</i>	5.01	4.89

*Table 1.* Predicted 2023 September extent and daily minimum extent using ice loss rates from different years or multi-year averages. Values in **bold** represent cases that would set new record lows.

This method yields a September 2023 extent of 4.77 ( $\pm 0.35$ ) million sq km. The large range in values, calculated from the standard deviation of the 16 years, is due to the limited number of years used and the significant amount of variability in daily extent change rates between 1 August and 30 September. Using the standard 30-year 1981-2010 climatology, the September extent is 5.01 million sq km. The lowest projected extent (from the last 18 years), which is from the 2012 rates, is 4.01 million sq km, while the highest, from 2006 rates, is 5.30 million square kilometers. The daily extent trajectories for the two averages and the high and low years from the last decade are provided in the figure below and the prediction using the rates for each of the last 18 years are provided in Table 1.

There isn't much expected skill at this point because of the large range of extent loss rates that may still occur. However, it provides a reasonable envelop of physically realistic September extents. As September approaches, the "window" of possible extents narrows and hones in on the final observed extent. An updated projection will plan to be submitted in September, using 2023 extent data through 1 September.




**Figure 1.** Predicted Arctic trajectories from the 1 August observed extent. The official estimate is based on the projection used the average of the last 16 years (2007-2022, green solid line) and for the 1981-2010 average (solid red line). The rates for 2006 and 2012 (dotted lines) yield the highest and lowest predicted extents respectively.

We use the same methodology for the Antarctic. However, since the trend has not particularly accelerated there isn't a reason to use any particular subset of years. It

is clear from Figure 2 that there is very little difference between using the thirty-year average versus the 16-year (2007-2022) average employed for the Arctic. So, the projection for the Antarctic is based on the daily rates of change for the 2007-2022 average to be consistent with the Arctic. The highest project results from using 2006 rates and the lowest projection comes from 2015 rates.

Using the 2007-2022 rates, the projected Antarctic September average is 16.22 ( $\pm 0.34$ ) million square kilometers. This represents an increase of  $\sim 0.5$  million square kilometers from the June predicting, indicative of the very slow increase in Antarctic sea ice extent during the month of June. Over the last 18 years, the highest September extent of 16.74 million square kilometers came from using 2022 rates, and the lowest September extent, using 2015 rates is 15.70 million square kilometers. The very large spread between the highest and lowest is not surprising given the high interannual variability in Antarctic sea ice. Since the Antarctic sea ice at its maximum encircles the entire continent over a thousand kilometers from the coast in most regions, even relatively small differences in ice edge location can result in large differences in total area. The maximum daily extent is predicted to be 16.30 ( $\pm 0.34$ ) million square kilometers and occurs on 26 September.



**Figure 2.** Predicted Antarctic trajectories from the 1 August observed extent. The official estimate is based on the projection used the average of the last 16 years (2007-2022, green solid line) and for the 1981-2010 average (solid red line). The rates for 2006 and 2015 (dotted lines) yield the highest and lowest predicted extents respectively.

### **Input Data Set References**

Maslanik, J. and J. Stroeve. 1999, updated daily. Near-Real-Time DMSP SSMIS Daily Polar Gridded Sea Ice Concentrations, Version 1. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center. <https://doi.org/10.5067/U8C09DWVX9LM>.

Fetterer, F., K. Knowles, W. Meier, M. Savoie, and A. K. Windnagel. 2017, updated daily. Sea Ice Index, Version 3. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center. <https://doi.org/10.7265/N5K072F8>.